



Organised systems of regular follow-up and review can improve blood pressure control

Clinical question

What interventions can improve control of blood pressure in patients with hypertension?

Bottom line

An organised system of registration, recall and regular review, allied to a vigorous stepped care approach to antihypertensive treatment, reduced blood pressure and all-cause mortality in a single, large randomised control trial. Health professional (nurse or pharmacist) led care appears to be a promising way of delivering care but requires further evaluation.

Caveat

Trials of educational interventions directed at patients or health professionals were heterogeneous and appear unlikely to be associated with large net reductions in blood pressure by themselves.

Context

Hypertension is a common problem in general practice. International community based studies show blood pressure goals are achieved in only 25–40% of patients who take antihypertensives.¹ There is a paucity of evidence as to how care for hypertensive patients should be delivered in the community to help improve blood pressure control.

Cochrane Systematic Review

Fahey T et al. Interventions used to improve control of blood pressure in patients with hypertension. Cochrane Database Syst Rev. 2006;(2):CD005182.

This review contains 56 trials with sizes ranging from 15 to 7772 participants.

PEARLS No. 269, September 2007, written by Brian R McAvoy

Further references

1. Burnier M. *Journal of Hypertension* 2002;20:1251–3

PEARLS are succinct summaries of Cochrane Systematic Reviews for primary care practitioners – developed by the Cochrane Primary Care Field, New Zealand Branch of the Australasian Cochrane Centre at the Department of General Practice and Primary Health Care, University of Auckland and funded by the New Zealand Guidelines Group. New Zealanders can access the Cochrane Library free via www.nzgg.org.nz

PEARLS provide guidance on whether a treatment is effective or ineffective. PEARLS are prepared as an educational resource and do not replace clinician judgement in the management of individual cases. View PEARLS online at: www.nzdoctor.co.nz; www.nzgg.org.nz; www.cochraneprimarycare.org